Prepared for submission to JHEP

arXiv:1303.5119v1 [hep-ph] 20 Mar 2013

Renormalization of the second-order QCD with arbitrary chromomagnetic factor and quark self-interactions.

Carlos A. Vaquera-Araujo

Facultad de Ciencias, CUICBAS, Universidad de Colima,
Bernal Díaz del Castillo 340, Colima, México, 28045.

E-mail: carolusvaquera@gmail.com

Abstract: In this work, the renormalization of the second-order QCD is analyzed at one-loop level, including an arbitrary chromomagnetic factor and quark self-interactions, which are mass dimension four operators in this framework. The model is shown to be renormalizable for any value of the chromomagnetic factor κ in the validity region of the perturbative expansion, and Dirac QCD is recovered in the $\kappa = 2$ case with vanishing self-interaction couplings.

Contents

1 Introduction **1**

2 The model **2**
2.1 Lagrangian 2
2.2 Feynman rules 3

3 Renormalization **6**
3.1 Counterterms 6
3.2 Superficial degree of divergence 7

4 Divergent amplitudes **8**
4.1 Gluon self-energy 8
4.2 Fermion self-energy 9
4.3 Ghost self-energy 10
4.4 Quark-quark-gluon vertex 10
4.5 Ghost-ghost-gluon vertex 11
4.6 Three-gluon vertex 12
4.7 Quark-quark-gluon-gluon vertex 12
4.8 Four-gluon vertex 14
4.9 Four-quark vertex 15

5 Beta functions **19**

6 Summary and conclusions **20**

1 Introduction

The idea of a second order formalism for spin 1/2 fermions can be traced back to Feynman [1] and Gell-Mann [2]. Applied to QCD, this formalism has proved to be very useful in the efficient determination of multi-gluon amplitudes due to its resemblance with scalar calculations (see for example [3] and further references). Important developments can be found in [4] and specially in [5] for the non-abelian gauge theory. More recently, an analysis of the anomalous chromomagnetic moment couplings of the top quark was performed in [6], where a second order formalism was used.

In general, an anomalous magnetic moment can be implemented naturally in a second order theory, and besides, it can be properly renormalized at one-loop level in the abelian version if point-like four-fermion interactions are included [9, 10]. Notice that this is possible because fermion fields have mass dimension one in this case. The purpose of this work is to generalize this result to a non-abelian gauge theory, analyzing the one-loop renormalization of the QCD of second order fermions with an arbitrary chromomagnetic factor in presence of fermion self-interactions. The starting point of this paper is the Poincaré-projector formalism [7, 8], a framework based on the projection onto irreducible representations of the Poincaré group, constructed to solve the problems of the quantum description of interacting high spin fields.

This paper is organized as follows: In Section 2, the basics of the model are presented. The renormalization procedure is sketched in Section 3. Section 4 is devoted to the calculation of the divergent piece of all the potentially divergent amplitudes of the theory and Section 5 to the determination of the corresponding beta functions. Finally, the conclusions of the work are discussed in Section 6.

2 The model

2.1 Lagrangian

The basic dimension four $SU(N_c)$ gauge invariant Lagrangian in the Poincaré-projector formalism is given by

$$\mathscr{L}_{SU(N_c)} = (\overline{D_\mu q}) K^{\mu\nu} (D_\nu q) - m^2 \bar{q} q - \frac{1}{4} G^{a\mu\nu} G^a_{\mu\nu}, \tag{2.1}$$

where q are the quark fields, living in the fundamental representation of $SU(N_c)$, with an internal $SU(N_f)$ global flavor symmetry (all flavors are degenerate with mass m) and covariant derivatives

$$D_\mu q = \left(\partial_\mu - i g A^a_\mu t^a\right) q. \tag{2.2}$$

A^a_μ stands for the gluon field, t^a are the generators of $SU(N_c)$ ($1 \leq a \leq N_c^2 - 1$), and the field strength tensor is

$$G^a_{\mu\nu} = \partial_\mu A^a_\nu - \partial_\nu A^a_\mu + g f^{abc} A^b_\mu A^c_\nu, \tag{2.3}$$

with structure constants f^{abc} defined by

$$[t^a, t^b] = i f^{abc}. \tag{2.4}$$

The space-time tensor $K^{\mu\nu}$ in the quark kinetic term is defined as

$$K^{\mu\nu} \equiv g^{\mu\nu} - i\kappa M^{\mu\nu} - i\kappa' \tilde{M}^{\mu\nu}, \tag{2.5}$$

with the Lorentz generators $M^{\mu\nu}$ for the $(1/2, 0) \oplus (0, 1/2)$ representation, their dual generators $\tilde{M}^{\mu\nu} = \epsilon^{\mu\nu\alpha\beta} M_{\alpha\beta}/2$ and arbitrary constants κ, κ'. Here κ can be identified with the chromomagnetic factor, and κ' parameterizes parity violating chromoelectric dipole interactions (see [9, 10] for further details and conventions). Setting $\kappa = 2$ and $\kappa' = 0$, eq. (2.1) describes the second order formalism for QCD [5]. In this paper, the analysis is restricted to $\kappa' = 0$, with κ kept arbitrary.

In order to obtain the gluon propagator from eq. (2.1), the gauge is fixed through a covariant gauge contribution and the Faddeev-Popov ghosts $\bar{c}^a$, c^a using

$$\mathscr{L}_{\text{gauge}} = -\frac{1}{2\xi} \left(\partial^\mu A^a_\mu\right)^2 + \left(\partial^\mu \bar{c}^a\right)\left(D_\mu c^a\right), \tag{2.6}$$

with the ghost covariant derivative: $D_\mu c^a = \left(\partial_\mu \delta^{ac} + g f^{abc} A^b_\mu\right) c^c$, where the unit matrix in the adjoint representation δ^{ab} is written down explicitly. Introducing an auxiliary scalar field B^a, this Lagrangian can be written, up to a total divergence, as

$$\mathscr{L}_{\text{gauge}} = -\frac{\xi}{2}(B^a)^2 + B^a \partial^\mu A^a_\mu - \bar{c}^a \left(\partial^\mu D_\mu c^a\right), \tag{2.7}$$

As stated in the introduction, second order fermion fields have mass dimension 1 in four space-time dimensions, and therefore, also Nambu-Jona-Lasinio-like interactions [11, 12] are dimension-four and gauge-invariant. The possible self-interactions (involving only elements of the $M^{\mu\nu}$ algebra) are:

$$\begin{aligned} \mathscr{L}_{\text{self}} &= \frac{\lambda_S}{2}\left(\bar{q}q\right)^2 + \frac{\lambda_P}{2}\left(\bar{q}\gamma^5 q\right)^2 + \frac{\lambda_T}{2}\left(\bar{q}M^{\mu\nu}q\right)^2 \\ &+ \frac{\lambda_{S_t}}{2}\left(\bar{q}t^a q\right)^2 + \frac{\lambda_{P_t}}{2}\left(\bar{q}\gamma^5 t^a q\right)^2 + \frac{\lambda_{T_t}}{2}\left(\bar{q}M^{\mu\nu}t^a q\right)^2 . \end{aligned} \tag{2.8}$$

Thus, the complete Lagrangian of the model is

$$\mathscr{L} = \mathscr{L}_{SU(N_c)} + \mathscr{L}_{\text{gauge}} + \mathscr{L}_{\text{self}}. \tag{2.9}$$

Notice that this Lagrangian is BRST invariant, as can be checked with the transformations

$$\begin{aligned} \delta A^a_\mu &= \theta D_\mu c^a, \\ \delta q &= i g \theta c^a t^a q, \\ \delta c^a &= -\frac{1}{2} g \theta f^{abc} c^b c^c, \\ \delta \bar{c}^a &= \theta B^a, \\ \delta B^a &= 0, \end{aligned} \tag{2.10}$$

parameterized by the infinitesimal anticommuting constant θ.

2.2 Feynman rules

The quadratic part of $\mathscr{L}$ gives the propagators shown in figure 1. Defining $\square\,[p] \equiv p^2 - m^2 + i\varepsilon$ and $\triangle\,[q] \equiv q^2 + i\varepsilon$, the quark propagator is

$$iS(p) = \frac{i}{\square\,[p]} \mathbb{1}_s \mathbb{1}_c \mathbb{1}_f, \tag{2.11}$$

with spinor unit matrix $\mathbb{1}_s$, color unit matrix $\mathbb{1}_c$ and flavor unit matrix $\mathbb{1}_f$. The gluon propagator is

$$iD^{ab}_{\mu\nu}(q) = -\frac{i}{\triangle\,[q]}\left[g_{\mu\nu} + (\xi - 1)\frac{q_\mu q_\nu}{q^2}\right]\delta^{ab}, \tag{2.12}$$

and the ghost propagator is

$$iG^{ab}(k) = \frac{i}{\triangle\,[k]}\delta^{ab}. \tag{2.13}$$



Figure 1. Propagators.

The Feynman rules for the interaction vertices of $\mathscr{L}$ are shown in figure 2, with the following definitions:

$$\begin{aligned} W^{\mu\nu\rho}(q_1,q_2,q_3) =& (q_1-q_2)^\rho g^{\mu\nu} + (q_2-q_3)^\mu g^{\nu\rho} + (q_3-q_1)^\nu g^{\rho\mu}, && (2.14)\\ W^{abcd}_{\mu\nu\rho\sigma} =& f^{abe}f^{cde}\left(g_{\mu\rho}g_{\nu\sigma}-g_{\mu\sigma}g_{\nu\rho}\right) + f^{ace}f^{bde}\left(g_{\mu\nu}g_{\rho\sigma}-g_{\mu\sigma}g_{\nu\rho}\right) && (2.15)\\ &+ f^{ade}f^{bce}\left(g_{\mu\nu}g_{\rho\sigma}-g_{\mu\rho}g_{\nu\sigma}\right), \\ V^\mu(p',p) =& \left[(p'+p)^\mu \mathbb{1}_s + i\kappa M^{\mu\nu}(p'-p)_\nu\right]\mathbb{1}_f, && (2.16)\\ V^{ab}_{\mu\nu} =& \Big[g_{\mu\nu}\{t^a,t^b\}\mathbb{1}_s - i\kappa M_{\mu\nu}[t^a,t^b]\Big]\mathbb{1}_f. && (2.17) \end{aligned}$$

The 4-quark vertex is given by

$$i\Lambda = i\left(\lambda_S \mathbb{1}\overline{\otimes}\mathbb{1} + \lambda_P \chi\overline{\otimes}\chi + \lambda_T M\overline{\otimes}M + \lambda_{S_t} t\overline{\otimes}t + \lambda_{P_t}\chi_t\overline{\otimes}\chi_t + \lambda_{T_t}M_t\overline{\otimes}M_t\right), \tag{2.18}$$

where the following notation is adopted

$$\begin{aligned} \mathbb{1}\overline{\otimes}\mathbb{1} &= \left[\delta_{i_1i_2}\delta_{i_3i_4}\delta_{j_1j_2}\delta_{j_3j_4}\delta_{\alpha_1\alpha_2}\delta_{\alpha_3\alpha_4} - \delta_{i_1i_4}\delta_{i_3i_2}\delta_{j_1j_4}\delta_{j_3j_2}\delta_{\alpha_1\alpha_4}\delta_{\alpha_3\alpha_2}\right], \\ \chi\overline{\otimes}\chi &= \left[\delta_{i_1i_2}\delta_{i_3i_4}\delta_{j_1j_2}\delta_{j_3j_4}\gamma^5_{\alpha_1\alpha_2}\gamma^5_{\alpha_3\alpha_4} - \delta_{i_1i_4}\delta_{i_3i_2}\delta_{j_1j_4}\delta_{j_3j_2}\gamma^5_{\alpha_1\alpha_4}\gamma^5_{\alpha_3\alpha_2}\right], \\ M\overline{\otimes}M &= \left[\delta_{i_1i_2}\delta_{i_3i_4}\delta_{j_1j_2}\delta_{j_3j_4}M^{\mu\nu}_{\alpha_1\alpha_2}M_{\mu\nu\alpha_3\alpha_4} - \delta_{i_1i_4}\delta_{i_3i_2}\delta_{j_1j_4}\delta_{j_3j_2}M^{\mu\nu}_{\alpha_1\alpha_4}M_{\mu\nu\alpha_3\alpha_2}\right], \\ t\overline{\otimes}t &= \left[\delta_{i_1i_2}\delta_{i_3i_4}t^a_{j_1j_2}t^a_{j_3j_4}\delta_{\alpha_1\alpha_2}\delta_{\alpha_3\alpha_4} - \delta_{i_1i_4}\delta_{i_3i_2}t^a_{j_1j_4}t^a_{j_3j_2}\delta_{\alpha_1\alpha_4}\delta_{\alpha_3\alpha_2}\right], \\ \chi_t\overline{\otimes}\chi_t &= \left[\delta_{i_1i_2}\delta_{i_3i_4}t^a_{j_1j_2}t^a_{j_3j_4}\gamma^5_{\alpha_1\alpha_2}\gamma^5_{\alpha_3\alpha_4} - \delta_{i_1i_4}\delta_{i_3i_2}t^a_{j_1j_4}t^a_{j_3j_2}\gamma^5_{\alpha_1\alpha_4}\gamma^5_{\alpha_3\alpha_2}\right], \\ M_t\overline{\otimes}M_t &= \left[\delta_{i_1i_2}\delta_{i_3i_4}t^a_{j_1j_2}t^a_{j_3j_4}M^{\mu\nu}_{\alpha_1\alpha_2}M_{\mu\nu\alpha_3\alpha_4} - \delta_{i_1i_4}\delta_{i_3i_2}t^a_{j_1j_4}t^a_{j_3j_2}M^{\mu\nu}_{\alpha_1\alpha_4}M_{\mu\nu\alpha_3\alpha_2}\right], \end{aligned} \tag{2.19}$$

for explicit flavor ($1 \leq i_k \leq N_f$), color ($1 \leq j_k \leq N_c$) and spinor ($1 \leq \alpha_k \leq 4$) indices of the quarks labeled with momenta p_k ($1 \leq k \leq 4$) in the corresponding diagram of figure 2. For closed quark loops, a factor of -1 given by Fermi statistics must be included, as well as the appropriate symmetry factors for closed gluon loops. In second order QCD with $\kappa = 2$, an additional $1/2$ factor can be attached to each spinor space trace in order to provide



Figure 2. Feynman rules for the second order QCD of self-interacting quarks with arbitrary chromomagnetic factor

a direct connection between the second order formalism and the Dirac one [4, 5]. In this work the convention used in [10] is adopted, where the trace of the spinor unit matrix is written as $\mathrm{tr}[\mathbb{1}_s] = \tau$, with $\tau = 4$ for the second order formalism in its natural form, and the formal relation $\tau = 2$ for comparison with Dirac QCD.

The $SU(N_c)$ conventions used in this work are

$$\begin{aligned} \mathrm{tr}[t^a t^b] &= T_F \delta^{ab}, \\ t^a t^a &= C_F, \qquad C_F = T_F \left(N_c - \frac{1}{N_c} \right), \\ if^{acd} if^{bdc} &= C_A \delta^{ab}, \qquad C_A = 2 T_F N_c, \end{aligned} \tag{2.20}$$

with T_F as the normalization constant for generator traces, and C_F (C_A) as the Casimir of the fundamental (adjoint) representation.

3 Renormalization

3.1 Counterterms

The free and interacting pieces of the bare Lagrangian are

$$\begin{aligned}\mathscr{L}_{\text{free}} =& -\frac{1}{4}\left(\partial^\mu A_0^{a\nu} - \partial^\nu A_0^{a\mu}\right)\left(\partial_\mu A_{0\nu}^a - \partial_\nu A_{0\mu}^a\right) - \frac{1}{2\xi_0}\left(\partial^\mu A_{0\mu}\right)^2 \\ &+ \left(\partial^\mu \bar{c}_0^a\right)\left(\partial_\mu c_0^a\right) + \left(\partial^\mu \bar{q}_0\right)\left(\partial_\mu q_0\right) - m_0^2 \bar{q}_0 q_0,\end{aligned} \tag{3.1}$$

$$\begin{aligned}\mathscr{L}_{\text{int}} =& -\frac{g}{2} f^{abc}\left(\partial^\mu A_0^{a\nu} - \partial^\nu A_0^{a\mu}\right) A_{0\mu}^b A_{0\nu}^c \\ &- \frac{g^2}{4} f^{eab} f^{ecd} A_{0\mu}^a A_{0\nu}^b A_0^{c\mu} A_0^{d\nu} - g f^{abc}\left(\partial_\mu \bar{c}_0^a\right) c_0^b A^{c\mu} \\ &+ i g_0\left[\bar{q}_0 K_{0\nu\mu}\left(\partial^\mu q_0\right) - \left(\partial^\mu \bar{\psi}_0\right) K_{0\mu\nu}\psi_0\right] A_0^{a\nu} t^a + g_0^2 \bar{q}_0 K_{0\mu\nu} q_0 A_0^{a\mu} t^a A_0^{b\nu} t^b \\ &+ \frac{\lambda_{S0}}{2}\left(\bar{q}_0 q_0\right)^2 + \frac{\lambda_{P0}}{2}\left(\bar{q}_0\gamma^5 q_0\right)^2 + \frac{\lambda_{T0}}{2}\left(\bar{q}_0 M^{\mu\nu} q_0\right)^2 \\ &+ \frac{\lambda_{S_t0}}{2}\left(\bar{q}_0 t^a q_0\right)^2 + \frac{\lambda_{P_t0}}{2}\left(\bar{q}_0\gamma^5 t^a q_0\right)^2 + \frac{\lambda_{T_t0}}{2}\left(\bar{q}_0 M^{\mu\nu} t^a q_0\right)^2.\end{aligned} \tag{3.2}$$

with $K_0^{\mu\nu} \equiv g^{\mu\nu} - i\kappa_0 M^{\mu\nu}$.

Introducing the renormalized fields

$$A_r^\mu = Z_A^{-\frac{1}{2}} A_0^\mu, \qquad q_r = Z_q^{-\frac{1}{2}} q_0, \qquad c_r = Z_c^{-\frac{1}{2}} c_0, \tag{3.3}$$

together with the renormalized parameters

$$\begin{aligned} g_r &= Z_g^{-1} g_0, & m_r &= Z_m^{-\frac{1}{2}} m_0, & \kappa_r &= Z_\kappa^{-1}\kappa_0, \\ \lambda_{Sr} &= Z_S^{-1}\lambda_{S0}, & \lambda_{Pr} &= Z_P^{-1}\lambda_{P0}, & \lambda_{Tr} &= Z_T^{-1}\lambda_{T0}, \\ \lambda_{S_tr} &= Z_{S_t}^{-1}\lambda_{S_t0}, & \lambda_{P_tr} &= Z_{P_t}^{-1}\lambda_{P_t0}, & \lambda_{T_tr} &= Z_{T_t}^{-1}\lambda_{T_t0}, \\ & & \xi_r &= Z_A^{-1}\xi_0, & & \end{aligned} \tag{3.4}$$

and the notation $Z \equiv 1 + \delta$, the free and interacting Lagrangians can be written as

$$\begin{aligned}\mathscr{L}_{\text{free}} =& \mathscr{L}_{\text{free}}^r - \frac{1}{4}\left(\partial^\mu A_r^{a\nu} - \partial^\nu A_r^{a\mu}\right)\left(\partial_\mu A_{r\nu}^a - \partial_\nu A_{r\mu}^a\right)\delta_A \\ &+ \left(\partial^\mu \bar{c}_r^a\right)\left(\partial_\mu c_r^a\right) + \left(\partial^\mu \bar{c}_r^a\right)\left(\partial_\mu c_r^a\right)\delta_c \\ &+ \left(\partial^\mu \bar{q}_r\right)\left(\partial_\mu q_r\right)\delta_q - m_r^2 \bar{q}_r q_r \delta_{mq^2},\end{aligned} \tag{3.5}$$

$$\begin{aligned}\mathscr{L}_{\text{int}} =& \mathscr{L}_{\text{int}}^r - \frac{g_r}{2} f^{abc}\left(\partial^\mu A_r^{a\nu} - \partial^\nu A_r^{a\mu}\right) A_{r\mu}^b A_{r\nu}^c \delta_{A^3} \\ &- \frac{g_r^2}{4} f^{eab} f^{ecd} A_{r\mu}^a A_{r\nu}^b A_r^{c\mu} A_{r\nu}^{d\nu}\delta_{A^4} - g_r f^{abc}\left(\partial_\mu \bar{c}_r^a\right) c_r^b A^{c\mu}\delta_{Ac^2} \\ &+ i g_r\left[\bar{q}_r g_{\nu\mu}\left(\partial^\mu q_r\right) - \left(\partial^\mu \bar{q}_r\right) g_{\mu\nu} q_r\right] A_r^{a\nu} t^a \delta_{Aq^2} \\ &+ i g_r\left[\bar{q}_r(-i\kappa_r M_{\mu\nu})\left(\partial^\mu q_r\right) - \left(\partial^\mu \bar{q}_r\right)(-i\kappa_r M_{\mu\nu}) q_r\right] A_r^{a\nu} t^a \delta_{\kappa A q^2} \\ &+ g_r^2 \bar{q}_r g_{\mu\nu} q_r A_r^{a\mu} t^a A_r^{b\nu} t^b \delta_{A^2 q^2} + g_r^2 \bar{q}_r(-i\kappa_r M_{\mu\nu}) q_r A_r^{a\mu} t^a A_r^{b\nu} t^b \delta_{\kappa A^2 q^2} \\ &\frac{\lambda_{Sr}}{2}\delta_{Sq^4}\left(\bar{q}_r q_r\right)^2 + \frac{\lambda_{Pr}}{2}\delta_{Pq^4}\left(\bar{q}_r\gamma^5 q_r\right)^2 + \frac{\lambda_{Tr}}{2}\delta_{Tq^4}\left(\bar{q}_r M^{\mu\nu} q_r\right)^2 \\ &+ \frac{\lambda_{S_tr}}{2}\delta_{S_tq^4}\left(\bar{q}_r t^a q_r\right)^2 + \frac{\lambda_{P_tr}}{2}\delta_{P_tq^4}\left(\bar{q}_r\gamma^5 t^a q_r\right)^2 + \frac{\lambda_{T_tr}}{2}\delta_{T_tq^4}\left(\bar{q}_r M^{\mu\nu} t^a q_r\right)^2,\end{aligned} \tag{3.6}$$

where $\mathscr{L}^r_{\text{free}}$ ($\mathscr{L}^r_{\text{int}}$) is the same Lagrangian as $\mathscr{L}_{\text{free}}$ ($\mathscr{L}_{\text{int}}$) with renormalized fields and couplings. The new renormalization constants introduced in eqs. (3.5) and (3.6) are defined as follows:

$$\begin{array}{rlrlrl}
Z_{mq^2} &= Z_m Z_q, & Z_{A^3} &= Z_g Z_A^{\frac{3}{2}}, & Z_{A^4} &= Z_g^2 Z_A^2, \\
Z_{Ac^2} &= Z_g Z_A^{\frac{1}{2}} Z_c, & Z_{Aq^2} &= Z_g Z_A^{\frac{1}{2}} Z_q, & Z_{A^2q^2} &= Z_g^2 Z_A Z_q, \\
Z_{\kappa A q^2} &= Z_\kappa Z_g Z_A^{\frac{1}{2}} Z_q, & Z_{\kappa A^2 q^2} &= Z_\kappa Z_g^2 Z_A Z_q, & Z_{Sq^4} &= Z_S Z_q^2, \\
Z_{Pq^4} &= Z_P Z_q^2, & Z_{Tq^4} &= Z_T Z_q^2, & Z_{S_t q^4} &= Z_{S_t} Z_q^2, \\
Z_{P_t q^4} &= Z_{P_t} Z_q^2, & Z_{T_t q^4} &= Z_{T_t} Z_q^2. & &
\end{array} \tag{3.7}$$

In the following, for simplicity, the suffix r will be removed in the renormalized parameters, keeping the suffix 0 for the bare quantities. In this notation, the Feynman rules for the renormalized fields are given in figures 1 and 2, while the corresponding rules for the counterterms are shown in figures 3 and 4.


ν
q
μ
b
a
$i(q^\mu q^\nu - g^{\mu\nu} q^2)\delta^{ab}\delta_A$
p
$i(p^2\delta_q - m^2\delta_{mq^2})\mathbb{1}_s\mathbb{1}_c\mathbb{1}_f$
k
b
a
$ik^2\delta^{ab}\delta_c$

Figure 3. Counterterms for Propagators.

3.2 Superficial degree of divergence

The superficial degree of divergence D for this theory is analogous to that of scalar QCD (SQCD). A Feynman diagram with N_G external gluon lines, N_F external quark lines and N_{FP} ghost lines has

$$D \leq 4 - N_G - N_F - \frac{3}{2}N_{FP}. \tag{3.8}$$

The factor 3/2 takes into account the fact that half of the ghost internal lines adjacent to ghost external lines in the Feynman diagram do not contribute to D due to the peculiar form of the ghost-ghost-gluon vertex.

The eight superficially divergent Feynman amplitudes are the gluon, ghost and quark self-energy parts and the 3-gluon, ghost-ghost-gluon, quark-quark-gluon, 4-gluon and quark-quark-gluon-gluon vertices. The renormalization of these vertices will be carried at one-loop level for arbitrary covariant gauge in dimensional regularization and in the $\overline{\text{MS}}$ scheme, using a commuting γ^5 with $M^{\mu\nu}$ in $d = 4 - 2\epsilon$ dimensions. As in [10], γ^5 is present only in diagrams with self-interactions, and therefore, pure QCD diagrams are free from


μ, a
q_1
q_2
q_3
ν, b
ρ, c
$g f^{abc} W^{\mu\nu\rho}(q_1, q_2, q_3)\delta_{A^3}$
q, μ, a
q, ν, b
p_1
p_2
$ig^2 g_{\mu\nu}\{t^a, t^b\}\mathbb{1}_s\mathbb{1}_f\delta_{A^2\psi^2}$
$+g^2\kappa M_{\mu\nu}[t^a, t^b]\mathbb{1}_f\delta_{\kappa A^2\psi^2}$
μ, a
p_2
p_1
$igt^a(p_1 + p_2)^\mu\delta_{Aq^2}$
$-gt^a\kappa M^{\mu\nu}(p_1 - p_2)_\nu\delta_{\kappa Aq^2}$
μ, a
ν, b
ρ, c
σ, d
$-ig^2 W^{abcd}_{\mu\nu\rho\sigma}\delta_{A^4}$
μ, b
k, c
k', a
$-g f^{abc} k'_\mu \delta_{Ac^2}$
p_2
p_1
p_4
p_3
$i\lambda_S\delta_{Sq^2}\mathbb{1}\overline{\otimes}\mathbb{1} + i\lambda_{S_t}\delta_{S_tq^2}t\overline{\otimes}t$
$+i\lambda_P\delta_{Pq^2}\chi\overline{\otimes}\chi + i\lambda_{P_t}\delta_{P_tq^2}\chi_t\overline{\otimes}\chi_t$
$+i\lambda_T\delta_{Tq^2}M\overline{\otimes}M + i\lambda_{T_t}\delta_{T_tq^2}M_t\overline{\otimes}M_t$

Figure 4. Counterterms for 3-point and 4-point vertices.

dimensional regularization inconsistencies. The same observations made in [10] regarding self-interactions are applicable in this work, and further study must be done in this direction.

In d dimensions, the renormalized parameters scale according to

$$\begin{array}{lll} g_r \to \mu^\epsilon g_r, & m_r \to m_r, & \kappa_r \to \kappa_r, \\ \lambda_{Sr} \to \mu^{2\epsilon}\lambda_{Sr}, & \lambda_{Pr} \to \mu^{2\epsilon}\lambda_{Pr}, & \lambda_{Tr} \to \mu^{2\epsilon}\lambda_{Tr}, \\ \lambda_{S_tr} \to \mu^{2\epsilon}\lambda_{S_tr}, & \lambda_{P_tr} \to \mu^{2\epsilon}\lambda_{P_tr}, & \lambda_{T_tr} \to \mu^{2\epsilon}\lambda_{T_tr}. \end{array} \tag{3.9}$$

In the next section, the divergent pieces of the superficially divergent amplitudes are extracted in order to determine the value of the renormalization constants at one-loop precision.

4 Divergent amplitudes

4.1 Gluon self-energy

The gluon self-energy at one-loop level is

$$-i\Pi^{ab}_{\mu\nu}(q) = -i\Pi^{*ab}_{\mu\nu}(q) - i\delta_A\left(g_{\mu\nu}q^2 - q_\mu q_\nu\right)\delta^{ab}, \tag{4.1}$$

where $-i\Pi^{*ab}_{\mu\nu}(q)$ stands for the contribution of the one-loop diagrams in Figure 5. As expected, $\Pi^{ab}_{\mu\nu}(q)$ is transverse

$$\Pi^{ab}_{\mu\nu}(q) = (g^{\mu\nu}q^2 - q^\mu q^\nu)\delta^{ab}\Pi(q^2), \tag{4.2}$$



Figure 5. Feynman diagrams for the gluon self-energy at one-loop.



Figure 6. Feynman diagrams for the fermion self-energy at one-loop.

and the divergent structure of the $\Pi(q^2)$ form factor is given by

$$\Pi(q^2) = \frac{g^2}{(4\pi)^2}\left[\tau N_f T_F \left(\frac{3\kappa^2-4}{12}\right) - C_A\left(\frac{13}{6}-\frac{\xi}{2}\right)\right]\frac{1}{\tilde{\epsilon}} + \delta_A + \text{finite}. \tag{4.3}$$

Thus, in the $\overline{\text{MS}}$ subtraction scheme, the gluon-field renormalization constant at one loop is

$$Z_A = 1 - \frac{g^2}{(4\pi)^2}\left[\tau N_f T_F \left(\frac{3\kappa^2-4}{12}\right) - C_A\left(\frac{13}{6}-\frac{\xi}{2}\right)\right]\frac{1}{\tilde{\epsilon}}. \tag{4.4}$$

4.2 Fermion self-energy

The fermion self-energy at one-loop level reads

$$-i\Sigma(p^2) = -i\Sigma^*(p^2) + ip^2\delta_q - im^2\delta_{mq^2}, \tag{4.5}$$

where $-i\Sigma^*(p^2)$ is computed with the one-loop diagrams of figure 6 and yields

$$\Sigma(p^2) = p^2\Sigma_1(p^2) + m^2\Sigma_2(p^2). \tag{4.6}$$



Figure 7. Feynman diagram for the ghost self-energy at one-loop.

where the form factors are

$$\Sigma_1(p^2) = \frac{g^2 C_F (3-\xi)}{(4\pi)^2} \frac{1}{\tilde{\epsilon}} - \delta_q + \text{finite}, \tag{4.7}$$

$$\begin{aligned} \Sigma_2(p^2) =& \frac{1}{(4\pi)^2} \Big\{ g^2 C_F \left(\xi + \frac{3\kappa^2}{4} \right) + C_F \left[\lambda_{S_t} + \lambda_{P_t} + 3\lambda_{T_t} \right] \\ & - \left[(\tau N_f N_C - 1)\lambda_S - \lambda_P - 3\lambda_T \right] \Big\} \frac{1}{\tilde{\epsilon}} + \delta_{mq^2} + \text{finite}. \end{aligned} \tag{4.8}$$

Form these results, the Z_q and Z_{mq^2} renormalization constants in the $\overline{\text{MS}}$ scheme are

$$Z_q = 1 + \frac{g^2 C_F (3-\xi)}{(4\pi)^2} \frac{1}{\tilde{\epsilon}}, \tag{4.9}$$

$$\begin{aligned} Z_{mq^2} =& 1 - \frac{1}{(4\pi)^2} \Big\{ g^2 C_F \left(\xi + \frac{3\kappa^2}{4} \right) + C_F \left[\lambda_{S_t} + \lambda_{P_t} + 3\lambda_{T_t} \right] \\ & - \left[(\tau N_f N_C - 1)\lambda_S - \lambda_P - 3\lambda_T \right] \Big\} \frac{1}{\tilde{\epsilon}}. \end{aligned} \tag{4.10}$$

4.3 Ghost self-energy

The ghost self-energy at one-loop level is

$$-i\widetilde{\Sigma}^{ab}(k^2) = -i\widetilde{\Sigma}^{*ab}(k^2) + ik^2 \delta^{ab} \delta_c, \tag{4.11}$$

where $\widetilde{\Sigma}^{*ab}(k^2)$ corresponds to the diagram of figure 7. A direct calculation gives

$$\widetilde{\Sigma}^{ab}(p^2) = \delta^{ab} k^2 \left[\frac{g^2 C_A}{(4\pi)^2} \left(\frac{3-\xi}{4} \right) \frac{1}{\tilde{\epsilon}} - \delta_c \right] + \text{finite}, \tag{4.12}$$

and the ghost-field $\overline{\text{MS}}$ renormalization constant is determined as

$$Z_c = 1 + \frac{g^2 C_A}{(4\pi)^2} \left(\frac{3-\xi}{4} \right) \frac{1}{\tilde{\epsilon}}. \tag{4.13}$$

4.4 Quark-quark-gluon vertex

From the Feynman rules, the qqg vertex at one-loop level is

$$\begin{aligned} ig\Gamma^{a\mu}(p_1, p_2, q) =& igt^a V^\mu(p_1, p_2) + ig\Gamma^{*a\mu}(p_1, p_2, q) \\ & + igt^a V^\mu(p_1, p_2)\delta_{Aq^2} + igt^a [i\kappa M^{\mu\nu}(p_1 - p_2)_\nu]\delta_{\kappa A q^2}, \end{aligned} \tag{4.14}$$



Figure 8. Feynman diagrams for the qqg vertex at one-loop.

where $ig\Gamma^{*a\mu}(p', p, q)$ stands for the contribution of the one-loop diagrams in figure 8. The divergent piece of this amplitude lives in two form factors

$$\Gamma^{a\mu}(p_1, p_2, q) = t^a(p_1 + p_2)^\mu \Gamma_1 + it^a \kappa M^{\mu\nu}(p_1 - p_2)_\nu \Gamma_2 + \text{finite}, \tag{4.15}$$

given by

$$\begin{aligned} \Gamma_1 =& -\frac{g^2}{(4\pi)^2}\left[C_F(3-\xi) - C_A\left(\frac{3+\xi}{4}\right)\right]\frac{1}{\tilde{\epsilon}} + \delta_{Aq^2} + \text{finite}, & (4.16)\\ \Gamma_2 =& \frac{1}{(4\pi)^2}\left\{\frac{g^2}{4}\left[\frac{C_A}{2}\left(\kappa^2 + 4\kappa - 6 + 2\xi\right) - C_F\left(\kappa^2 + 8 - 4\xi\right)\right] + \lambda_S + \lambda_P - \lambda_T \right. & (4.17)\\ & \left. + \left(C_F - \frac{C_A}{2}\right)\left[\lambda_{S_t} + \lambda_{P_t} - \lambda_{T_t}\right] + \frac{\tau}{2} N_f T_F \lambda_{T_t}\right\}\frac{1}{\tilde{\epsilon}} + \delta_{\kappa A q^2} + \text{finite}. \end{aligned}$$

Therefore, the Z_{Aq^2} and $Z_{\kappa A q^2}$ renormalization constants in the $\overline{\text{MS}}$ scheme are

$$\begin{aligned} Z_{Aq^2} =& 1 + \frac{g^2}{(4\pi)^2}\left[C_F(3-\xi) - C_A\left(\frac{3+\xi}{4}\right)\right]\frac{1}{\tilde{\epsilon}}, & (4.18)\\ Z_{\kappa A q^2} =& 1 - \frac{1}{(4\pi)^2}\left\{\frac{g^2}{4}\left[\frac{C_A}{2}\left(\kappa^2 + 4\kappa - 6 + 2\xi\right) - C_F\left(\kappa^2 + 8 - 4\xi\right)\right]\right. & (4.19)\\ & + \lambda_S + \lambda_P - \lambda_T + \frac{\tau}{2} N_f T_F \lambda_{T_t} \\ & \left. + \left(C_F - \frac{C_A}{2}\right)\left[\lambda_{S_t} + \lambda_{P_t} - \lambda_{T_t}\right]\right\}\frac{1}{\tilde{\epsilon}}. \end{aligned}$$

4.5 Ghost-ghost-gluon vertex

The ccg vertex at one-loop level is unmodified by the second order formalism for fermions

$$-g\widetilde{\Gamma}^{abc\mu}(k_1, k_2, q) = -gf^{abc}k_1^\mu - g\widetilde{\Gamma}^{*abc\mu}(k_1, k_2, q) - gf^{abc}k_1^\mu \delta_{Ac^2}, \tag{4.20}$$



Figure 9. Feynman diagrams for the ccg vertex at one-loop.

with $-g\widetilde{\Gamma}^{*abc\mu}(k',k,q)$ calculated from the one-loop diagrams in figure 9. The divergent part of this amplitude is

$$\widetilde{\Gamma}^{abc\mu}(k',k,q) = f^{abc}k_1^{\mu}\left[\frac{g^2}{(4\pi)^2}\frac{C_A}{2}\xi\frac{1}{\tilde{\epsilon}} + \delta_{Ac^2}\right] + \text{finite}, \tag{4.21}$$

and the Z_{Ac^2} $\overline{\text{MS}}$ renormalization constant reads

$$Z_{Ac^2} = 1 - \frac{g^2}{(4\pi)^2}\frac{C_A}{2}\xi\frac{1}{\tilde{\epsilon}}. \tag{4.22}$$

4.6 Three-gluon vertex

The ggg vertex at one-loop level is

$$g\mathcal{W}^{abc}_{\mu\nu\rho}(q_1,q_2,q_3) = gf^{abc}W_{\mu\nu\rho}(q_1,q_2,q_3) + g\mathcal{W}^{*abc}_{\mu\nu\rho}(q_1,q_2,q_3) + gf^{abc}W_{\mu\nu\rho}(q_1,q_2,q_3)\delta_{A^3}, \tag{4.23}$$

where $g\mathcal{W}^{*abc}_{\mu\nu\rho}(q_1,q_2,q_3)$ is the contribution of the one-loop diagrams in figure 10. The divergent piece of this amplitude is contained in one form factor

$$\mathcal{W}^{abc}_{\mu\nu\rho}(q_1,q_2,q_3) = f^{abc}W_{\mu\nu\rho}(q_1,q_2,q_3)\mathcal{W} + \text{finite}, \tag{4.24}$$

given by

$$\mathcal{W} = \frac{g^2}{(4\pi)^2}\left[\tau N_f T_F\left(\frac{3\kappa^2-4}{12}\right) - C_A\left(\frac{17}{12} - \frac{3\xi}{4}\right)\right]\frac{1}{\tilde{\epsilon}} + \delta_{A^3} + \text{finite}, \tag{4.25}$$

which, in turn, determines the Z_{A^3} renormalization constant in the $\overline{\text{MS}}$ scheme as

$$Z_{A^3} = 1 - \frac{g^2}{(4\pi)^2}\left[\tau N_f T_F\left(\frac{3\kappa^2-4}{12}\right) - C_A\left(\frac{17}{12} - \frac{3\xi}{4}\right)\right]\frac{1}{\tilde{\epsilon}}. \tag{4.26}$$

4.7 Quark-quark-gluon-gluon vertex

The $qqgg$ vertex function at one-loop level is obtained from the Feynman rules as

$$\begin{aligned} ig^2\mathcal{G}^{ab}_{\mu\nu}(p_1,p_2,q_1,q_2) =& ig^2V^{ab}_{\mu\nu} + ig^2\mathcal{G}^{*ab}_{\mu\nu}(p_1,p_2,q_1,q_2) \\ &+ ig^2g_{\mu\nu}\{t^a,t^b\}\delta_{A^2q^2} - ig^2\left[i\kappa M_{\mu\nu}\right][t^a,t^b]\delta_{\kappa A^2q^2}, \end{aligned} \tag{4.27}$$



Figure 10. Feynman diagrams for the ggg vertex at one-loop.

where the one-loop corrections $ig^2\mathcal{G}^{*ab}_{\mu\nu}(p_1,p_2,q_1,q_2)$ are given by the diagrams of figure 11. As expected, the divergent pieces of this amplitude are contained in two form factors

$$\mathcal{G}^{ab}_{\mu\nu}(p_1,p_2,q_1,q_2) = g_{\mu\nu}\{t^a,t^b\}\mathcal{G}_1 - i\kappa M_{\mu\nu}[t^a,t^b]\mathcal{G}_2 + \text{finite}, \tag{4.28}$$

read as

$$\mathcal{G}_1 = -\frac{g^2}{(4\pi)^2}\left[C_F(3-\xi) - C_A\left(\frac{3+\xi}{2}\right)\right]\frac{1}{\tilde{\epsilon}} + \delta_{A^2q^2} + \text{finite}, \tag{4.29}$$

$$\begin{aligned}\mathcal{G}_2 =&\frac{1}{(4\pi)^2}\Bigg\{\frac{g^2}{4}\left[\frac{C_A}{2}\left(\kappa^2+4\kappa+4\xi\right) - C_F\left(\kappa^2+8-4\xi\right)\right] + \lambda_S + \lambda_P - \lambda_T \\ &+ \left(C_F - \frac{C_A}{2}\right)\left[\lambda_{S_t} + \lambda_{P_t} - \lambda_{T_t}\right] + \frac{\tau}{2}N_f T_F \lambda_{T_t}\Bigg\}\frac{1}{\tilde{\epsilon}} + \delta_{\kappa A^2 q^2} + \text{finite}.\end{aligned} \tag{4.30}$$



Figure 11. Feynman diagrams for the $qqgg$ vertex at one-loop.

From these results, the $Z_{A^2q^2}$ and $Z_{\kappa A^2q^2}$ renormalization constants in the $\overline{\text{MS}}$ scheme are given by

$$
\begin{aligned}
Z_{A^2q^2} =& 1 + \frac{g^2}{(4\pi)^2}\left[C_F(3-\xi) - C_A\left(\frac{3+\xi}{2}\right)\right]\frac{1}{\tilde{\epsilon}}, && (4.31)\\
Z_{\kappa A^2q^2} =& 1 - \frac{1}{(4\pi)^2}\left\{\frac{g^2}{4}\left[\frac{C_A}{2}\left(\kappa^2+4\kappa+4\xi\right) - C_F\left(\kappa^2+8-4\xi\right)\right]\right. && (4.32)\\
& \qquad + \lambda_S + \lambda_P - \lambda_T + \frac{\tau}{2}N_f T_F \lambda_{T_t}\\
& \qquad \left. + \left(C_F - \frac{C_A}{2}\right)\left[\lambda_{S_t} + \lambda_{P_t} - \lambda_{T_t}\right]\right\}\frac{1}{\tilde{\epsilon}}.
\end{aligned}
$$

4.8 Four-gluon vertex

The $gggg$ vertex at one-loop level is

$$
-ig^2\mathcal{E}^{abcd}_{\mu\nu\rho\sigma}(q_1,q_2,q_3,q_4) = -\,ig^2W^{abcd}_{\mu\nu\rho\sigma} - ig^2\mathcal{E}^{*abcd}_{\mu\nu\rho\sigma}(q_1,q_2,q_3,q_4) - ig^2W^{abcd}_{\mu\nu\rho\sigma}\delta_{A^4}, \quad (4.33)
$$



Figure 12. Basic Feynman diagrams for the $gggg$ vertex at one-loop. The number indicates the total number of diagrams obtained from each basic form under permutation of external gluon lines.

where $-ig^2\mathcal{E}^{abcd}_{\mu\nu\rho\sigma}(q_1,q_2,q_3,q_4)$ is the contribution of the one-loop diagrams in figure 12. The divergent piece of this amplitude is contained in one form factor

$$\mathcal{E}^{abcd}_{\mu\nu\rho\sigma}(q_1,q_2,q_3,q_4) = W^{abcd}_{\mu\nu\rho\sigma}\mathcal{E} + \text{finite}, \tag{4.34}$$

given by

$$\mathcal{E} =\frac{g^2}{(4\pi)^2}\left[\tau N_f T_F\left(\frac{3\kappa^2-4}{12}\right) - C_A\left(\frac{2}{3}-\xi\right)\right]\frac{1}{\tilde{\epsilon}} + \delta_{A^4} + \text{finite}, \tag{4.35}$$

and the corresponding $\overline{\text{MS}}$ renormalization constant yields

$$Z_{A^4} =1 - \frac{g^2}{(4\pi)^2}\left[\tau N_f T_F\left(\frac{3\kappa^2-4}{12}\right) - C_A\left(\frac{2}{3}-\xi\right)\right]\frac{1}{\tilde{\epsilon}}. \tag{4.36}$$

4.9 Four-quark vertex

The last potentially divergent vertex, is the $qqqq$ vertex function

$$\begin{aligned} i\mathcal{F}\left(p_1,p_2,p_1',p_2'\right) &= i\Lambda + i\mathcal{F}^*\left(p_1,p_2,p_1',p_2'\right) + i\lambda_S\delta_{Sq^2}\mathbb{1}\overline{\otimes}\mathbb{1} + i\lambda_{S_t}\delta_{S_tq^2}t\overline{\otimes}t \\ &+ i\lambda_P\delta_{Pq^2}\chi\overline{\otimes}\chi + i\lambda_{P_t}\delta_{P_tq^2}\chi_t\overline{\otimes}\chi_t + i\lambda_T\delta_{Tq^2}M\overline{\otimes}M + i\lambda_{T_t}\delta_{T_tq^2}M_t\overline{\otimes}M_t, \end{aligned} \tag{4.37}$$

with $i\mathcal{F}^*\left(p_1,p_2,p_1',p_2'\right)$ obtained from the loop diagrams in figure 13. The divergent piece of this amplitude lives in six form factors

$$\begin{aligned} \mathcal{F}\left(p_1,p_2,p_1',p_2'\right) =&\mathcal{F}_1\mathbb{1}\overline{\otimes}\mathbb{1} + \mathcal{F}_2 t\overline{\otimes}t + \mathcal{F}_3\chi\overline{\otimes}\chi + \mathcal{F}_4\chi_t\overline{\otimes}\chi_t \\ &+ \mathcal{F}_5 M\overline{\otimes}M + \mathcal{F}_6 M_t\overline{\otimes}M_t + \text{finite}, \end{aligned} \tag{4.38}$$



Figure 13. Feynman diagrams for the $qqqq$ vertex function at one-loop.

given by

$$\begin{aligned}\mathcal{F}_1 =&\frac{1}{(4\pi)^2}\Bigg\{C_F\left(C_A-2C_F\right)\left[3g^4\left(1-\frac{\kappa^2}{4}\right)^2+\frac{3g^2\kappa^2}{4}\lambda_{T_t}+\lambda_{S_t}^2+\lambda_{P_t}^2+\frac{3}{2}\lambda_{T_t}^2\right] \\ &+2C_F\lambda_S\left(2g^2\xi+\frac{3g^2\kappa^2}{4}+\lambda_{S_t}+\lambda_{P_t}+3\lambda_{T_t}\right) \\ &+\lambda_S\left[\lambda_S(4-\tau N_f N_c)+2\lambda_P+6\lambda_T\right]+2\lambda_P^2+3\lambda_T^2\Bigg\}\frac{1}{\tilde{\epsilon}}+\lambda_S\delta_{Sq^2}+\text{finite},\end{aligned} \tag{4.39}$$

$$\begin{aligned}\mathcal{F}_2 =&\frac{1}{(4\pi)^2}\Bigg\{ \left(4C_F-\frac{3}{2}C_A\right)\left[3g^4\left(1-\frac{\kappa^2}{4}\right)^2+\frac{3g^2\kappa^2}{4}\lambda_{T_t}+\lambda_{S_t}^2+\lambda_{P_t}^2+\frac{3}{2}\lambda_{T_t}^2\right]\\ &+\lambda_{S_t}\left[2C_Fg^2\xi+(2C_F-C_A)\left(\frac{3g^2\kappa^2}{4}+\lambda_{S_t}+\lambda_{P_t}+3\lambda_{T_t}\right)\right]\\ &+\lambda_{S_t}\left[6(\lambda_S+\lambda_T)+2\lambda_P-\tau N_fT_F\lambda_{S_t}\right]\\ &+\lambda_T\left(\frac{3g^2\kappa^2}{2}+6\lambda_{T_t}\right)+4\lambda_P\lambda_{P_t}\Bigg\}\frac{1}{\tilde{\epsilon}}+\lambda_{S_t}\delta_{S_tq^2}+\text{finite},\end{aligned}\tag{4.40}$$

$$\begin{aligned}\mathcal{F}_3 =&\frac{1}{(4\pi)^2}\Bigg\{C_F\left(C_A-2C_F\right)\left[\frac{3g^2\kappa^2}{4}\lambda_{T_t}+2\lambda_{P_t}\lambda_{S_t}+\frac{3}{2}\lambda_{T_t}^2\right]\\ &+2C_F\lambda_P\left(2g^2\xi+\frac{3g^2\kappa^2}{4}+\lambda_{S_t}+\lambda_{P_t}+3\lambda_{T_t}\right)\\ &+\lambda_P\left[\lambda_P(2-\tau N_fN_c)+6(\lambda_S+\lambda_T)\right]+3\lambda_T^2\Bigg\}\frac{1}{\tilde{\epsilon}}+\lambda_P\delta_{Pq^2}+\text{finite},\end{aligned}\tag{4.41}$$

$$\begin{aligned}\mathcal{F}_4 =&\frac{1}{(4\pi)^2}\Bigg\{ \left(4C_F-\frac{3}{2}C_A\right)\left[\frac{3g^2\kappa^2}{4}\lambda_{T_t}+2\lambda_{S_t}\lambda_{P_t}+\frac{3}{2}\lambda_{T_t}^2\right]\\ &+\lambda_{P_t}\left[2C_Fg^2\xi+(2C_F-C_A)\left(\frac{3g^2\kappa^2}{4}+\lambda_{S_t}+\lambda_{P_t}+3\lambda_{T_t}\right)\right]\\ &+\lambda_{P_t}\left[6(\lambda_S+\lambda_T)+2\lambda_P-\tau N_fT_F\lambda_{P_t}\right]\\ &+\lambda_T\left(\frac{3g^2\kappa^2}{2}+6\lambda_{T_t}\right)+4\lambda_P\lambda_{S_t}\Bigg\}\frac{1}{\tilde{\epsilon}}+\lambda_{P_t}\delta_{P_tq^2}+\text{finite},\end{aligned}\tag{4.42}$$

$$\begin{aligned}\mathcal{F}_5 =&\frac{1}{(4\pi)^2}\Bigg\{C_F\left(C_A-2C_F\right)\left(\frac{g^2\kappa^2}{2}+2\lambda_{T_t}\right)\left(\lambda_{S_t}+\lambda_{P_t}\right)\\ &+2C_F\lambda_T\left(2g^2\xi-\frac{g^2\kappa^2}{4}+\lambda_{P_t}+\lambda_{S_t}-\lambda_{T_t}\right)\\ &+\lambda_T\left[6(\lambda_S+\lambda_P)-2\lambda_T\left(1+\frac{\tau}{4}N_fN_c\right)\right]\Bigg\}\frac{1}{\tilde{\epsilon}}+\lambda_T\delta_{Tq^2}+\text{finite},\end{aligned}\tag{4.43}$$

$$\begin{aligned}\mathcal{F}_6 =&\frac{1}{(4\pi)^2}\Bigg\{C_Ag^2\left[\frac{g^2\kappa^2}{4}\left(1-\frac{\kappa}{2}\right)^2-\frac{3\kappa^2}{4}\left(\lambda_{S_t}+\lambda_{P_t}-\lambda_{T_t}\right)\right]+g^2\kappa^2\left(\lambda_S+\lambda_P\right)\\ &+C_Fg^2\left[2\lambda_{T_t}\xi+\frac{\kappa^2}{2}\left(4\lambda_{S_t}+4\lambda_{P_t}-\lambda_{T_t}\right)\right]+4\lambda_T(\lambda_{S_t}+\lambda_{P_t})\\ &+2C_A\lambda_{T_t}\left(\lambda_{T_t}-2\lambda_{S_t}-2\lambda_{P_t}\right)+2C_F\lambda_{T_t}\left(5\lambda_{S_t}+5\lambda_{P_t}-\lambda_{T_t}\right)\\ &+\lambda_{T_t}\left[6\left(\lambda_S+\lambda_P\right)-2\lambda_T-\frac{\tau}{2}N_fT_F\lambda_{T_t}\right]\Bigg\}\frac{1}{\tilde{\epsilon}}+\lambda_{T_t}\delta_{T_tq^2}+\text{finite}.\end{aligned}\tag{4.44}$$

Finally, the $\overline{\mathrm{MS}}$ renormalization constants associated to this vertex are determined as

$$\begin{aligned}
Z_{Sq^2} = & 1 - \frac{1}{(4\pi)^2}\left\{ 2C_F\left(2g^2\xi + \frac{3g^2\kappa^2}{4} + \lambda_{S_t} + \lambda_{P_t} + 3\lambda_{T_t}\right)\right. \\
& + C_F\left(C_A - 2C_F\right)\frac{g^2}{\lambda_S}\left[3g^2\left(1 - \frac{\kappa^2}{4}\right)^2 + \frac{3\kappa^2}{4}\lambda_{T_t}\right] \\
& + C_F\left(C_A - 2C_F\right)\frac{1}{\lambda_S}\left(\lambda_{S_t}^2 + \lambda_{P_t}^2 + \frac{3}{2}\lambda_{T_t}^2\right) \\
& \left. + \lambda_S(4 - \tau N_f N_c) + 2\lambda_P + 6\lambda_T + 2\frac{\lambda_P^2}{\lambda_S} + 3\frac{\lambda_T^2}{\lambda_S}\right\}\frac{1}{\tilde{\epsilon}},
\end{aligned}\tag{4.45}$$

$$\begin{aligned}
Z_{S_tq^2} = & 1 - \frac{1}{(4\pi)^2}\left\{ 2C_F g^2\xi + (2C_F - C_A)\left(\frac{3g^2\kappa^2}{4} + \lambda_{S_t} + \lambda_{P_t} + 3\lambda_{T_t}\right)\right. \\
& + \left(4C_F - \frac{3}{2}C_A\right)\frac{g^2}{\lambda_{S_t}}\left[3g^2\left(1 - \frac{\kappa^2}{4}\right)^2 + \frac{3\kappa^2}{4}\lambda_{T_t}\right] \\
& + \left(4C_F - \frac{3}{2}C_A\right)\frac{1}{\lambda_{S_t}}\left(\lambda_{S_t}^2 + \lambda_{P_t}^2 + \frac{3}{2}\lambda_{T_t}^2\right) \\
& + 6(\lambda_S + \lambda_T) + 2\lambda_P - \tau N_f T_F \lambda_{S_t} \\
& \left. + \left(\frac{3g^2\kappa^2}{2} + 6\lambda_{T_t}\right)\frac{\lambda_T}{\lambda_{S_t}} + 4\lambda_P\frac{\lambda_{P_t}}{\lambda_{S_t}}\right\}\frac{1}{\tilde{\epsilon}},
\end{aligned}\tag{4.46}$$

$$\begin{aligned}
Z_{Pq^2} = & 1 - \frac{1}{(4\pi)^2}\left\{ C_F\left(C_A - 2C_F\right)\frac{1}{\lambda_P}\left[\frac{3g^2\kappa^2}{4}\lambda_{T_t} + 2\lambda_{P_t}\lambda_{S_t} + \frac{3}{2}\lambda_{T_t}^2\right]\right. \\
& + 2C_F\left(2g^2\xi + \frac{3g^2\kappa^2}{4} + \lambda_{S_t} + \lambda_{P_t} + 3\lambda_{T_t}\right) \\
& \left. + \lambda_P(2 - \tau N_f N_c) + 6(\lambda_S + \lambda_T) + 3\frac{\lambda_T^2}{\lambda_P}\right\}\frac{1}{\tilde{\epsilon}},
\end{aligned}\tag{4.47}$$

$$\begin{aligned}
Z_{P_tq^2} = & 1 - \frac{1}{(4\pi)^2}\left\{ \left(4C_F - \frac{3}{2}C_A\right)\frac{1}{\lambda_{P_t}}\left[\frac{3g^2\kappa^2}{4}\lambda_{T_t} + 2\lambda_{S_t}\lambda_{P_t} + \frac{3}{2}\lambda_{T_t}^2\right]\right. \\
& + 2C_F g^2\xi + (2C_F - C_A)\left(\frac{3g^2\kappa^2}{4} + \lambda_{S_t} + \lambda_{P_t} + 3\lambda_{T_t}\right) \\
& + 6(\lambda_S + \lambda_T) + 2\lambda_P - \tau N_f T_F \lambda_{P_t} \\
& \left. + \left(\frac{3g^2\kappa^2}{2} + 6\lambda_{T_t}\right)\frac{\lambda_T}{\lambda_{P_t}} + 4\lambda_P\frac{\lambda_{S_t}}{\lambda_{P_t}}\right\}\frac{1}{\tilde{\epsilon}},
\end{aligned}\tag{4.48}$$

$$\begin{aligned}
Z_{Tq^2} = & 1 - \frac{1}{(4\pi)^2}\left\{ C_F\left(C_A - 2C_F\right)\frac{1}{\lambda_T}\left(\frac{g^2\kappa^2}{2} + 2\lambda_{T_t}\right)(\lambda_{S_t} + \lambda_{P_t})\right. \\
& + 2C_F\left(2g^2\xi - \frac{g^2\kappa^2}{4} + \lambda_{P_t} + \lambda_{S_t} - \lambda_{T_t}\right) \\
& \left. + 6(\lambda_S + \lambda_P) - 2\lambda_T\left(1 + \frac{\tau}{4}N_f N_c\right)\right\}\frac{1}{\tilde{\epsilon}} + \lambda_T + \text{finite},
\end{aligned}\tag{4.49}$$

$$
\begin{aligned}
Z_{T_t q^2} =& 1 - \frac{1}{(4\pi)^2}\Bigg\{C_A g^2\left[\frac{g^2\kappa^2}{4\lambda_{T_t}}\left(1-\frac{\kappa}{2}\right)^2 - \frac{3\kappa^2}{4\lambda_{T_t}}\left(\lambda_{S_t}+\lambda_{P_t}-\lambda_{T_t}\right)\right] \\
&+ C_F g^2\left[2\xi + \frac{\kappa^2}{2\lambda_{T_t}}\left(4\lambda_{S_t}+4\lambda_{P_t}-\lambda_{T_t}\right)\right] + \frac{g^2\kappa^2}{\lambda_{T_t}}\left(\lambda_S+\lambda_P\right) \\
&+ 2C_A\left(\lambda_{T_t}-2\lambda_{S_t}-2\lambda_{P_t}\right) + 2C_F\left(5\lambda_{S_t}+5\lambda_{P_t}-\lambda_{T_t}\right) \\
&+ 6\left(\lambda_S+\lambda_P\right) - 2\lambda_T - \frac{\tau}{2}N_f T_F \lambda_{T_t} + 4\frac{\lambda_T}{\lambda_{T_t}}(\lambda_{S_t}+\lambda_{P_t})\Bigg\}\frac{1}{\tilde{\epsilon}}.
\end{aligned}
\tag{4.50}
$$

5 Beta functions

Form the renormalization constants obtained in eqs.(4.4), (4.9), (4.10), (4.13), (4.18), (4.19), (4.31), (4.32), (4.22), (4.26), (4.36), (4.45), (4.46), (4.47), (4.48), (4.49), (4.50) and their definition in eq. (3.7), it can be shown that the following Slavnov-Taylor identity hold at one loop:

$$
\frac{Z_{Aq^2}}{Z_q} = \frac{Z_{A^2q^2}}{Z_{Aq^2}} = \frac{Z_{\kappa A^2 q^2}}{Z_{\kappa A q^2}} = \frac{Z_{Ac^2}}{Z_c} = \frac{Z_{A^3}}{Z_A} = \frac{Z_{A^3}}{Z_A} = \frac{Z_{A^4}}{Z_{A^3}} = 1 - \frac{g^2}{(4\pi)^2}C_A\left(\frac{3+\xi}{4}\right)\frac{1}{\tilde{\epsilon}}. \tag{5.1}
$$

The beta functions $\beta_\eta \equiv \mu\frac{\partial\eta}{\partial\mu}$ and anomalous dimensions $\gamma_m \equiv \frac{\mu}{m}\frac{\partial m}{\partial\mu}$ in the $\epsilon \to 0$ limit can be extracted from the obtained renormalization constants as

$$
\beta_g = -\frac{g^3}{(4\pi)^2}\left[\frac{11}{3}C_A - \tau N_f T_F\left(\frac{3\kappa^2-4}{12}\right)\right], \tag{5.2}
$$

$$
\begin{aligned}
\beta_\kappa =& \frac{\kappa}{(4\pi)^2}\Bigg\{\frac{g^2}{4}\left[2C_F(\kappa^2-4) - C_A(\kappa-2)(\kappa+6)\right] - \tau N_f T_F \lambda_{T_t} \\
&+ (C_A-2C_F)(\lambda_{S_t}+\lambda_{P_t}-\lambda_{T_t}) - 2(\lambda_S+\lambda_P-\lambda_T)\Bigg\},
\end{aligned}
\tag{5.3}
$$

$$
\begin{aligned}
\beta_{\lambda_S} =& -\frac{1}{(4\pi)^2}\Bigg\{\frac{3g^2}{8}\left[g^2\left(\kappa^2-4\right)^2 + 4\kappa^2\lambda_{T_t}\right]C_F(C_A-2C_F) \\
&+ C_F(C_A-2C_F)\left(2\lambda_{S_t}^2+2\lambda_{P_t}^2+3\lambda_{T_t}^2\right) \\
&+ C_F\lambda_S\left[3g^2\left(\kappa^2+4\right) + 4(\lambda_{P_t}+\lambda_{S_t}+3\lambda_{T_t})\right] \\
&+ 2\left[2\lambda_P^2 + 2\lambda_P\lambda_S + \lambda_S^2(4-\tau N_c N_F) + 6\lambda_S\lambda_T + 3\lambda_T^2\right]\Bigg\},
\end{aligned}
\tag{5.4}
$$

$$
\begin{aligned}
\beta_{\lambda_{S_t}} =& -\frac{1}{(4\pi)^2}\Bigg\{\frac{3g^2}{8}\left[g^2\left(\kappa^2-4\right)^2 + 4\kappa^2\lambda_{T_t} + 32\lambda_{S_t}\right]\left(4C_F - \frac{3}{2}C_A\right) \\
&+ \left(4C_F - \frac{3}{2}C_A\right)\left(2\lambda_{S_t}^2+2\lambda_{P_t}^2+3\lambda_{T_t}^2\right) + 3g^2\kappa^2\lambda_T \\
&+ \lambda_{S_t}\left(C_F - \frac{1}{2}C_A\right)\left[3g^2\left(\kappa^2-12\right) + 4(\lambda_{P_t}+\lambda_{S_t}+3\lambda_{T_t})\right] \\
&+ 2\left[2\lambda_P(2\lambda_{P_t}+\lambda_{S_t}) + 6\lambda_S\lambda_{S_t} - \tau N_f T_F \lambda_{S_t}^2 + 6\lambda_T(\lambda_{S_t}+\lambda_{T_t})\right]\Bigg\},
\end{aligned}
\tag{5.5}
$$

$$
\begin{aligned}
\beta_{\lambda_P} = -\frac{1}{(4\pi)^2}\Big\{ & C_F\left(C_A-2C_F\right)\left(\frac{3g^2\kappa^2}{2}\lambda_{T_t}+4\lambda_{P_t}\lambda_{S_t}+3\lambda_{T_t}^2\right) \\
& + C_F\lambda_P\left[3g^2\left(\kappa^2+4\right)+4(\lambda_{P_t}+\lambda_{S_t}+3\lambda_{T_t})\right] \\
& + 2\left[\lambda_P^2(2-\tau N_c N_F)+6\lambda_P(\lambda_S+\lambda_T)+3\lambda_T^2\right]\Big\},
\end{aligned} \tag{5.6}
$$

$$
\begin{aligned}
\beta_{\lambda_{P_t}} = -\frac{1}{(4\pi)^2}\Big\{ & \left(4C_F-\frac{3}{2}C_A\right)\left[\frac{3g^2}{2}\left(8\lambda_{P_t}+\kappa^2\lambda_{T_t}\right)+4\lambda_{P_t}\lambda_{S_t}+3\lambda_{T_t}^2\right] \\
& + \left(C_F-\frac{1}{2}C_A\right)\lambda_{P_t}\left[3g^2\left(\kappa^2-12\right)+4(\lambda_{P_t}+\lambda_{S_t}+3\lambda_{T_t})\right] \\
& + 2\left[2\lambda_P(\lambda_{P_t}+2\lambda_{S_t})-\tau N_f T_F\lambda_{P_t}^2+6\lambda_{P_t}\lambda_S+6\lambda_T(\lambda_{P_t}+\lambda_{T_t})\right] \\
& + 3g^2\kappa^2\lambda_T\Big\},
\end{aligned} \tag{5.7}
$$

$$
\begin{aligned}
\beta_{\lambda_T} = -\frac{1}{(4\pi)^2}\Big\{ & C_F(C_A-2C_F)(\lambda_{P_t}+\lambda_{S_t})\left(g^2\kappa^2+4\lambda_{T_t}\right) \\
& + C_F\lambda_T\left[g^2(12-\kappa^2)+4(\lambda_{P_t}+\lambda_{S_t}-\lambda_{T_t})\right] \\
& + \lambda_T\left[12\lambda_P+12\lambda_S-\lambda_T(\tau N_c N_f+4)\right]\Big\},
\end{aligned} \tag{5.8}
$$

$$
\begin{aligned}
\beta_{\lambda_{T_t}} = -\frac{1}{(4\pi)^2}\Big\{ & C_A g^2\left[\frac{1}{8}g^2(\kappa-2)^2\kappa^2-\frac{3\kappa^2}{2}(\lambda_{P_t}+\lambda_{S_t}-\lambda_{T_t})\right] \\
& + 4C_A\lambda_{T_t}\left[\lambda_{T_t}-2(\lambda_{P_t}+\lambda_{S_t})\right]+C_F g^2\kappa^2\left[4(\lambda_{P_t}+\lambda_{S_t})-\lambda_{T_t}\right] \\
& + 4C_F\lambda_{T_t}\left[3g^2+5(\lambda_{P_t}+\lambda_{S_t})-\lambda_{T_t}\right]+2(\lambda_P+\lambda_S)\left(g^2\kappa^2+6\lambda_{T_t}\right) \\
& + 8\lambda_T(\lambda_{P_t}+\lambda_{S_t})-4\lambda_T\lambda_{T_t}-\lambda_{T_t}^2\tau N_f T_F\Big\},
\end{aligned} \tag{5.9}
$$

$$
\begin{aligned}
\gamma_m = -\frac{1}{(4\pi)^2}\Big\{ & C_F\left[\frac{3g^2}{4}\left(\kappa^2+4\right)+(\lambda_{P_t}+\lambda_{S_t}+3\lambda_{T_t})\right] \\
& + (\lambda_P-\tau N_c N_f\lambda_S+\lambda_S+3\lambda_T)\Big\}.
\end{aligned} \tag{5.10}
$$

Notice that there is a fixed point for eqs. (5.3)–(5.9) at $\kappa = 2$ with all self-interaction couplings set to zero. In this fixed point, the theory has a simple connection to Dirac QCD, which amounts to take formally $\tau \to 2$ (See [10] for an analogous discussion in QED). This is the only choice available for pure second order QCD (vanishing self-interaction couplings).

6 Summary and conclusions

In this work, the one-loop level renormalization of QCD in the second order Poincaré projector formalism was performed. The description of an arbitrary chromomagnetic factor is natural in this framework and fermion self-interactions can be incorporated without spoiling renormalizability. As expected, the conclusions of this analysis are analogous to those of [10], and can be summarized as follows: *i*) Pure second order QCD is renormalizable only for $\kappa = 2$. *ii*) Fermion self-interactions are also renormalizable at one-loop and make possible the renormalization of an arbitrary chromomagnetic factor in the second order formalism. Finally it is important to remark that the last conclusion requires a deeper analysis due to the γ^5 issues of the dimensional regularization method.

Acknowledgments

This work was supported by CONACyT under project CB-2011-167425. The author acknowledges Mauro Napsuciale and Alfredo Aranda for their helpful comments and critical reading of the manuscript.

References

[1] R. P. Feynman, *An Operator calculus having applications in quantum electrodynamics*, *Phys. Rev.* **84** (1951) 108.

[2] R. P. Feynman and M. Gell-Mann, *Theory of Fermi interaction*, *Phys. Rev.* **109** (1958) 193.

[3] C. Schubert, *Perturbative quantum field theory in the string inspired formalism*, *Phys. Rept.* **355** (2001) 73 [hep-th/0101036].

[4] M. J. G. Veltman, *Two component theory and electron magnetic moment*, *Acta Phys. Polon.* **B 29** (1998) 783 [hep-th/9712216].

[5] A. G. Morgan, *Second order fermions in gauge theories*, *Phys. Lett.* **B 351** (1995) 249 [hep-ph/9502230].

[6] A. J. Larkoski and M. E. Peskin, Phys. Rev. D **83**, 034012 (2011) [arXiv:1012.0552 [hep-ph]].

[7] E. G. Delgado-Acosta, M. Napsuciale, S. Rodriguez, *Second order formalism for spin 1/2 fermions and Compton scattering*, *Phys. Rev.* **D 83** (2011) 073001 [arXiv:1012.4130 [hep-ph]].

[8] M. Napsuciale, M. Kirchbach, S. Rodriguez, *Spin 3/2 beyond the Rarita-Schwinger framework*, *Eur. Phys. J.* **A 29** (2006) 289 [hep-ph/0606308].

[9] R. Angeles-Martinez and M. Napsuciale, *Renormalization of the QED of second order spin 1/2 fermions*, *Phys. Rev.* **D 85** (2012) 076004 [arXiv:1112.1134].

[10] C. A. Vaquera-Araujo, M. Napsuciale and R. Angeles-Martinez, *Renormalization of the QED of Self-Interacting Second Order Spin 1/2 Fermions*, JHEP **1301**, 011 (2013) [arXiv:1205.1557 [hep-ph]].

[11] Y. Nambu and G. Jona-Lasinio, *Dynamical model of elementary particles based on an analogy with superconductivity. I*, *Phys. Rev.* **122** (1961) 345.

[12] Y. Nambu and G. Jona-Lasinio, *Dynamical model of elementary particles based on an analogy with superconductivity. II*, *Phys. Rev.* **124** (1961) 246.